

## NOTCH1 Protein, Mouse (HEK293, His)

Cat. No.:	HY-P71166
Synonyms:	Neurogenic locus notch homolog protein 1; Notch 1; Motch A; mT14
Species:	Mouse
Source:	HEK293
Accession:	Q01705 (A18-Q526)
Gene ID:	18128
Molecular Weight:	Approximately 70.0 kDa

### PROPERTIES

AA Sequence	<pre> A A R G L R C S Q P   S G T C L N G G R C   E V A N G T E A C V   C S G A F V G Q R C Q D S N P C L S T P   C K N A G T C H V V   D H G G T V D Y A C   S C P L G F S G P L C L T P L D N A C L   A N P C R N G G T C   D L L T L T E Y K C   R C P P G W S G K S C Q Q A D P C A S N   P C A N G G Q C L P   F E S S Y I C R C P   P G F H G P T C R Q D V N E C S Q N P G   L C R H G G T C H N   E I G S Y R C A C R   A T H T G P H C E L P Y V P C S P S P C   Q N G G T C R P T G   D T T H E C A C L P   G F A G Q N C E E N V D D C P G N N C K   N G G A C V D G V N   T Y N C R C P P E W   T G Q Y C T E D V D E C Q L M P N A C Q   N G G T C H N T H G   G Y N C V C V N G W   T G E D C S E N I D D C A S A A C F Q G   A T C H D R V A S F   Y C E C P H G R T G   L L C H L N D A C I S N P C N E G S N C   D T N P V N G K A I   C T C P S G Y T G P   A C S Q D V D E C A L G A N P C E H A G   K C L N T L G S F E   C Q C L Q G Y T G P   R C E I D V N E C I S N P C Q N D A T C   L D Q I G E F Q C I   C M P G Y E G V Y C   E I N T D E C A S S P C L H N G H C M D   K I N E F Q C Q C P   K G F N G H L C Q </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

## Background

NOTCH1 Protein functions as a receptor for membrane-bound ligands, including Jagged-1 (JAG1), Jagged-2 (JAG2), and Delta-1 (DLL1), playing a crucial role in cell-fate determination. Upon ligand activation, the released Notch intracellular domain (NICD) forms a transcriptional activator complex with RBPJ/RBPSUH, activating genes in the enhancer of split locus and influencing differentiation, proliferation, and apoptosis. NOTCH1 is involved in angiogenesis, negatively regulating endothelial cell proliferation, migration, and angiogenic sprouting. It contributes to thymic maturation of CD4(+) and CD8(+) cells, follicular differentiation, and possibly cell fate selection within the follicle. In cerebellar development, it serves as a receptor for neuronal DNER, impacting the differentiation of Bergmann glia. NOTCH1 also plays roles in repressing neuronal and myogenic differentiation, postimplantation development, mesoderm development, somite formation, and neurogenesis. Additionally, it modulates the balance between motile and immotile cilia for left/right symmetry determination. The protein interacts with various partners, including DNER, DTX1, DTX2, RBPJ/RBPSUH, MAML1, MAML2, MAML3, SNW1, AAK1, FBXW7, SGK1, HIF1AN, SNAI1, MDM2A, BCL6, THBS4, CCN3, DLL4, ZMIZ1, MEGF10, DLL1, JAG1, PRAG1, PSEN1, and ZFP64, highlighting its involvement in intricate signaling networks and regulatory processes.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA