

## Resistin Protein, Human (His)

<b>Cat. No.:</b>	HY-P71259
<b>Synonyms:</b>	Resistin; Adipose tissue-specific secretory factor; Cysteine-rich secreted protein FIZZ3; C/EBP-epsilon-regulated myeloid-specific secreted cysteine-rich protein; Cysteine-rich secreted protein A12-alpha-like 2; FIZZ3; HXCP1; RSTN; RETN
<b>Species:</b>	Human
<b>Source:</b>	E. coli
<b>Accession:</b>	Q9HD89 (K19-P108)
<b>Gene ID:</b>	56729
<b>Molecular Weight:</b>	10-14 kDa

### PROPERTIES

<b>AA Sequence</b>	K T L C S M E E A I      N E R I Q E V A G S      L I F R A I S S I G      L E C Q S V T S R G D L A T C P R G F A      V T G C T C G S A C      G S W D V R A E T T      C H C Q C A G M D W T G A R C C R V Q P
<b>Biological Activity</b>	Data is not available.
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20 mM Acetic acid, pH 3.0.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
<b>Storage &amp; Stability</b>	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA