

## ROR2 Protein, Human (HEK293, C-His)

<b>Cat. No.:</b>	HY-P71263A
<b>Synonyms:</b>	Tyrosine-protein kinase transmembrane receptor ROR2; Neurotrophic tyrosine kinase, receptor-related 2; ROR2; NTRKR2
<b>Species:</b>	Human
<b>Source:</b>	HEK293
<b>Accession:</b>	Q01974 (E34-G403)
<b>Gene ID:</b>	4920
<b>Molecular Weight:</b>	Approximately 55 kDa

### PROPERTIES

<b>AA Sequence</b>	<pre> E V E V L D P N D P   L G P L D G Q D G P   I P T L K G Y F L N   F L E P V N N I T I V Q G Q T A I L H C   K V A G N P P P N V   R W L K N D A P V V   Q E P R R I I I R K T E Y G S R L R I Q   D L D T T D T G Y Y   Q C V A T N G M K T   I T A T G V L F V R L G P T H S P N H N   F Q D D Y H E D G F   C Q P Y R G I A C A   R F I G N R T I Y V D S L Q M Q G E I E   N R I T A A F T M I   G T S T H L S D Q C   S Q F A I P S F C H F V F P L C D A R S   R T P K P R E L C R   D E C E V L E S D L   C R Q E Y T I A R S N P L I L M R L Q L   P K C E A L P M P E   S P D A A N C M R I   G I P A E R L G R Y H Q C Y N G S G M D   Y R G T A S T T K S   G H Q C Q P W A L Q   H P H S H H L S S T D F P E L G G G H A   Y C R N P G G Q M E   G P W C F T Q N K N   V R M E L C D V P S C S P R D S S K M G </pre>
<b>Biological Activity</b>	Measured in a cell inhibition assay by using A375 cells. The ED <sub>50</sub> this effect is 1.493 µg/mL, corresponding to a specific activity is 669.7924 units/mg.
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCL, 300 mM NaCl, pH 7.4.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
<b>Storage &amp; Stability</b>	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

---

**Background**

ROR2 protein, a tyrosine-protein kinase receptor, emerges as a key player in the early stages of chondrocyte formation and is deemed essential for cartilage and growth plate development. Its phosphorylation of YWHAB not only induces osteogenesis but also promotes bone formation. Despite exhibiting minimal tyrosine kinase activity in vitro, ROR2's intricate role extends to potentially acting as a receptor for the Wnt ligand WNT5A. This interaction may lead to the intriguing outcome of inhibiting WNT3A-mediated signaling. The multifaceted functions of ROR2 highlight its significance in orchestrating complex processes such as chondrogenesis, osteogenesis, and the intricate regulation of Wnt signaling pathways.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA