

Product Data Sheet

Serum Albumin/ALB Protein, Cynomolgus (P.pastoris, His)

Cat. No.: HY-P71711

Synonyms: ALB; Qlv-U347A-C12; Qlv-U377A-G8Serum albumin

Species: Cynomolgus Source: P. pastoris

Accession: A2V9Z4 (25D-608A)

Gene ID: 102130757

Molecular Weight: Approximately 68.0 kDa

PROPERTIES

AA Sequence	DTHKSEVAHR FKDLGEEHFK GLVLVAFSQY LQQCPFEEHV KLVNEVTEFA KTCVADESAE NCDKSLHTLF GDKLCTVATL RETYGEMADC CAKQEPERNE CFLQHKDDNP NLPPLVRPEV DVMCTAFHDN EATFLKKYLY EVARRHPYFY APELLFFAAR YKAAFAECCQ AADKAACLLP KLDELRDEGK ASSAKQRLKC ASLQKFGDRA FKAWAVARLS QKFPKAEFAE VSKLVTDLTK VHTECCHGDL LECADDRADL AKYMCENQDS ISSKLKECCD KPLLEKSHCL AEVENDEMPA DLPSLAADYV ESKDVCKNYA EAKDVFLGMF LYEYARRHPD YSVMLLLRLA KAYEATLEKC CAAADPHECY AKVFDEFQPL VEEPQNLVKQ NCELFEQLGE YKFQNALLVR YTKKVPQVST PTLVEVSRNL GKVGAKCCKL PEAKRMPCAE DYLSVVLNRL CVLHEKTPVS EKVTKCCTES LVNRRPCFSA LELDEAYVPK AFNAETFTFH ADMCTLSEKE KQVKKQTALV ELVKHKPKAT KEQLKGVMDN FAAFVEKCCK
Appearance	Lyophilized powder.
Formulation	Lyophilized after extensive dialysis against solution in Tris-based buffer, 50% glycerol.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Page 1 of 2 www. Med Chem Express. com

DESCRIPTION

Background

Serum Albumin (ALB) protein serves various roles, including binding water, calcium, sodium, potassium, fatty acids, hormones, bilirubin, and drugs. Its primary function involves regulating the colloidal osmotic pressure of blood. ALB acts as a major transporter of zinc in plasma, binding approximately 80% of all plasma zinc. It also serves as a significant transporter for calcium and magnesium, binding approximately 45% of circulating calcium and magnesium in plasma. ALB potentially possesses multiple calcium-binding sites and may additionally bind calcium in a non-specific manner. The presence of a shared binding site between zinc and calcium at residue Asp-273 suggests a potential interaction and communication between zinc and calcium transport in the blood, with zinc exhibiting higher affinity followed by calcium and magnesium. ALB also interacts with bacterial siderophore enterobactin, impeding enterobactin-mediated iron uptake by E.coli from ferric transferrin and thereby limiting the utilization of iron and the growth of enteric bacteria like E.coli. However, it does not hinder iron uptake by the bacterial siderophore aerobactin. ALB interacts with FCGRT to regulate its homeostasis and also interacts with TASOR. In plasma, ALB forms a covalently-linked complex with chromophore-bound alpha-1-microglobulin, yet this interaction does not hinder the binding of fatty acids to ALB.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA