

## TIM-3/HAOCR2 Protein, Mouse (HEK 293, hFc)

Cat. No.:	HY-P71995
Synonyms:	Havcr2; Tim3; Timd3; Hepatitis A virus cellular receptor 2 homolog; HAVcr-2; T-cell immunoglobulin and mucin domain-containing protein 3; TIMD-3; T-cell immunoglobulin mucin receptor 3; TIM-3; T-cell membrane protein 3; CD antigen CD366
Species:	Mouse
Source:	HEK 293
Accession:	Q8VIM0 (R20-R191)
Gene ID:	171285
Molecular Weight:	Approximately 50.0 kDa

### PROPERTIES

AA Sequence	<pre> R S L E N A Y V F E   V G K N A Y L P C S   Y T L S T P G A L V   P M C W G K G F C P W S Q C T N E L L R   T D E R N V T Y Q K   S S R Y Q L K G D L   N K G D V S L I I K N V T L D D H G T Y   C C R I Q F P G L M   N D K K L E L K L D   I K A A K V T P A Q T A H G D S T T A S   P R T L T T E R N G   S E T Q T L V T L H   N N N G T K I S T W A D E I K D S G E T   I R           </pre>
Biological Activity	The ED50 as determined by its ability to bind Human Galectin 9 in functional ELISA is less than 20 µg/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm solution of 1xPBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in sterile distilled water.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA