

SARS-CoV-2 S glycoprotein (HEK293, His-Myc)

Cat. No.:	HY-P72039
Synonyms:	S glycoprotein; E2; Peplomer protein
Species:	Virus
Source:	HEK293
Accession:	P0DTC2 (R319-F541)
Gene ID:	43740568
Molecular Weight:	Approximately 30.1 kDa

PROPERTIES

AA Sequence	<pre> RVQPTESIVR FPNITNLCPF GEVFNATRFA SVYAWNRRKI SNCVADYSVL YNSASFSTFK CYGVSPTKLN DLCFTNVYAD SFVIRGDEV R QIAPGQTGKI ADYNYKLPDD FTGCVIAWNS NNLDSKVGGN YNYLYRLFRK SNLKPFERDI STEIYQAGST PCNGVEGFNC YFPLQSYGFQ PTNGVGYQPY RVVVLSELL HAPATVCGPK KSTNLVKKNC VNF </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm solution of 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The SARS-CoV-2 S glycoprotein plays a crucial role in infection by attaching the virion to the host cell membrane through interaction with the primary receptor, host ACE2. Upon cleavage of S₂/S₂'₁, binding to the ACE2 receptor initiates either direct fusion at the cell membrane or internalization of the virus via endocytosis, leading to fusion of the virion membrane with the host endosomal membrane. Additionally, the glycoprotein may utilize NRP1/NRP2 and integrin as alternative entry receptors, possibly explaining the virus's tropism in human olfactory epithelial cells. The stalk domain of S exhibits three hinges, providing unexpected orientational freedom to the head. Acting as a class I viral fusion protein, the glycoprotein undergoes an extensive and irreversible conformational change during virus entry, triggered by host TMPRSS2 or CSTL,</p>
-------------------	--

leading to fusion of the viral envelope with the cellular cytoplasmic membrane and release of viral genomic RNA into the host cell cytoplasm. The glycoprotein exhibits at least three conformational states: pre-fusion native, pre-hairpin intermediate, and post-fusion hairpin, with the coiled coil regions adopting a trimer-of-hairpins structure during fusion, facilitating the apposition and subsequent fusion of viral and target cell membranes.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA