

OGG1 Protein, Human (N-His)

Cat. No.:	HY-P72338A
Synonyms:	8-oxoguanine DNA glycosylase; 8-hydroxyguanine DNA glycosylase; MMH; MUTM; OGH1
Species:	Human
Source:	E. coli
Accession:	O15527-1 (P2-G345)
Gene ID:	4968
Molecular Weight:	Approximately 38.7 kDa

PROPERTIES

AA Sequence	<p> P A R A L L P R R M G H R T L A S T P A L W A S I P C P R S E L R L D L V L P S G Q S F R W R E Q S P A H W S G V L A D Q V W T L T Q T E E Q L H C T V Y R G D K S Q A S R P T P D E L E A V R K Y F Q L D V T L A Q L Y H H W G S V D S H F Q E V A Q K F Q G V R L L R Q D P I E C L F S F I C S S N N N I A R I T G M V E R L C Q A F G P R L I Q L D D V T Y H G F P S L Q A L A G P E V E A H L R K L G L G Y R A R Y V S A S A R A I L E E Q G G L A W L Q Q L R E S S Y E E A H K A L C I L P G V G T K V A D C I C L M A L D K P Q A V P V D V H M W H I A Q R D Y S W H P T T S Q A K G P S P Q T N K E L G N F F R S L W G P Y A G W A Q A V L F S A D L R Q S R H A Q E P P A K R R K G S K G P E G </p>
Biological Activity	Data is not available.
Appearance	Solution
Formulation	Supplied as a 0.2 µm filtered solution of PBS, pH 7.4, 40% Glycerol.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice

DESCRIPTION

Background	OGG1 (8-Oxoguanine DNA Glycosylase 1) is a DNA repair enzyme crucial for maintaining genomic integrity. Specializing in
-------------------	---

the repair of oxidative DNA damage, OGG1 specifically incises DNA at 8-oxoG (8-oxoguanine) residues. Additionally, it excises other oxidatively damaged bases, such as 7,8-dihydro-8-oxoguanine and 2,6-diamino-4-hydroxy-5-N-methylformamidopyrimidine (FAPY), contributing to the removal of mutagenic lesions from the DNA. Notably, OGG1 possesses a beta-lyase activity, enabling it to nick the DNA strand 3' to the lesion, initiating the repair process. This enzymatic versatility highlights OGG1's critical role in safeguarding the genome by counteracting the detrimental effects of oxidative stress on DNA integrity.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA