

Product Data Sheet

CD24 Protein, Human (Biotinylated, HEK293, Fc-Avi)

| Cat. No.: | HY-P72352 |
|-------------------|--|
| Synonyms: | Signal transducer CD24; Small cell lung carcinoma cluster 4 antigen; CD24; CD24A |
| Species: | Human |
| Source: | HEK293 |
| Accession: | P25063 (S27-G59) |
| Gene ID: | 100133941 |
| Molecular Weight: | 40-55 kDa |
| | |

| PROPERTIES | |
|---------------------|--|
| AA Sequence | SETTTGTSSN SSQSTSNSGL APNPTNATTK AAG |
| Appearance | Lyophilized powder. |
| Formulation | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. |
| Endotoxin Level | <1 EU/µg, determined by LAL method. |
| Reconsititution | It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose). |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage. |
| Shipping | Room temperature in continental US; may vary elsewhere. |

DESCRIPTION

| Background | The CD24 protein is implicated in playing a pivotal role in the cell differentiation of various cell types, with its signaling potentially triggered by the binding of a lectin-like ligand to CD24 carbohydrates, transduced through the release of second messengers derived from the GPI-anchor. CD24 also functions in modulating B-cell activation responses, promoting AG-dependent proliferation of B-cells, and preventing their terminal differentiation into antibody-forming cells. In association with SIGLEC10, CD24 may contribute to the selective suppression of the immune response to danger-associated molecular patterns (DAMPs) such as HMGB1, HSP70, and HSP90, suggesting a role in immune regulation. Additionally, CD24 is implicated in the control of autoimmunity. It exhibits cell-type-specific interactions with LYN and FGR, binding to FRG in a |
|------------|--|
| | small cell cancer line, LYN in the erythroleukemia cell line K-562, and in Burkitt's lymphoma, highlighting its diverse involvement in cellular processes and immune regulation. |

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA