

## CD47 Protein, Human (Biotinylated, HEK293, Avi-His)

Cat. No.:	HY-P72359
Synonyms:	Leukocyte Surface Antigen CD47; Antigenic Surface Determinant Protein OA3; IAP; MER6
Species:	Human
Source:	HEK293
Accession:	Q08722 (Q19-P139)
Gene ID:	961
Molecular Weight:	30-55 kDa

### PROPERTIES

AA Sequence	<p>           Q L L F N K T K S V    E F T F C N D T V V    I P C F V T N M E A    Q N T T E V Y V K W            K F K G R D I Y T F    D G A L N K S T V P    T D F S S A K I E V    S Q L L K G D A S L            K M D K S D A V S H    T G N Y T C E V T E    L T R E G E T I I E    L K Y R V V S W F S            P         </p>
Biological Activity	<p>1. Immobilized Biotinylated Human CD47-His at 2 µg/mL (100 µl/well). Dose response curve for Human SIRP alpha-hFc with the EC<sub>50</sub> of 0.03 ng/mL determined by ELISA.</p> <p>2. Immobilized Human SIRP alpha, hFc Tag at 1 µg/mL (100 µl/well) on the plate. Dose response curve for Biotinylated Human CD47, His Tag with the EC<sub>50</sub> of &lt; 0.48 µg/mL determined by ELISA</p>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 10 mM Tris-Citrate, 150 mM NaCl, pH 8.0 or PBS (pH 7.4), normally 5% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	CD47, an adhesive protein, facilitates cell-to-cell interactions and serves as a receptor for thrombospondin THBS1, modulating integrin signaling through the activation of heterotrimeric G proteins. Involved in diverse cellular processes,
------------	--

---

CD47 contributes to signal transduction, cardiovascular homeostasis, inflammation, apoptosis, angiogenesis, cellular self-renewal, and immunoregulation. Notably, it plays a role in modulating pulmonary endothelin EDN1 signaling and functions as a pressor agent in the regulation of blood pressure in response to THBS1. CD47 is crucial for memory formation and synaptic plasticity in the hippocampus, acting as a receptor for SIRPA and SIRPG, which impacts dendritic cell maturation, cytokine production, cell-cell adhesion, and T-cell activation. Furthermore, CD47 positively modulates FAS-dependent apoptosis in T-cells and suppresses angiogenesis, contributing to metabolic dysregulation during aging. In response to THBS1, CD47 negatively modulates wound healing, inhibits stem cell self-renewal, and may play a role in membrane transport and/or integrin-dependent signal transduction. As a monomer, CD47 interacts with THBS1, SIRPA, FAS/CD95, SIRPG, UBQLN1, UBQLN2, and potentially fibrinogen, highlighting its intricate involvement in cellular and molecular pathways.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA