

## TIM-3/HAVCR2 Protein, Cynomolgus (HEK293, His)

<b>Cat. No.:</b>	HY-P72447
<b>Synonyms:</b>	Hepatitis A virus cellular receptor 2 homolog; T cell immunoglobulin and mucin domain3; HAVCR2; CD366; TIM3
<b>Species:</b>	Cynomolgus
<b>Source:</b>	HEK293
<b>Accession:</b>	G7P6Q7 (S22-R201)
<b>Gene ID:</b>	10214172
<b>Molecular Weight:</b>	26-30 kDa

### PROPERTIES

<b>AA Sequence</b>	<p>           S E V E Y I A E V G    Q N A Y L P C S Y T    P A P P G N L V P V    C W G K G A C P V F            D C S N V V L R T D    N R D V N D R T S G    R Y W L K G D F H K    G D V S L T I E N V            T L A D S G V Y C C    R I Q I P G I M N D    E K H N V K L V V I    K P A K V T P A P T            L Q R D L T S A F P    R M L T T G E H G P    A E T Q T P G S L P    D V N L T V S N F F            C E L Q I F T L T N    E L R D S G A T I R         </p>
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
<b>Storage &amp; Stability</b>	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

<b>Background</b>	<p>TIM-3/HAVCR2 protein, a cell surface receptor, plays a pivotal role in modulating both innate and adaptive immune responses. Predominantly considered as having an inhibitory function, the reported stimulating functions suggest a nuanced influence based on cellular context and ligand specificity. It regulates macrophage activation and inhibits T-helper type 1 lymphocyte (Th1)-mediated auto- and alloimmune responses, promoting immunological tolerance. In CD8+ cells, TIM-3 attenuates TCR-induced signaling by blocking NF-kappaB and NFAT promoter activities, leading to reduced IL-2 secretion. This inhibitory function is proposed to involve its association with LCK, impairing the phosphorylation of TCR subunits. Conversely, TIM-3 has been shown to activate TCR-induced signaling in T-cells, likely implicating ZAP70, LCP2,</p>
-------------------	---

---

LCK, and FYN. The receptor for LGALS9, TIM-3's binding to LGALS9 is believed to suppress T-cell responses, leading to apoptosis of antigen-specific cells. Additionally, TIM-3 may recognize phosphatidylserine on apoptotic cells, mediating their phagocytosis, and positively regulates innate immune responses, particularly in dendritic cells. It also plays a role in suppressing nucleic acid-mediated innate immune responses in tumor-infiltrating dendritic cells and negatively regulates NK cell function in LPS-induced endotoxic shock. Interactions with various signaling molecules, including HMGB1, BAG6, PIK3R1, PIK3R2, FYN, and ILF3, further contribute to the intricate regulatory functions of TIM-3 in immune responses.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA