

## Siglec-15 Protein, Human (HEK293, His)

Cat. No.:	HY-P72475
Synonyms:	Sialic acid-binding Ig-like lectin 15; Siglec-15; CD33 antigen-like 3; CD33L3
Species:	Human
Source:	HEK293
Accession:	Q6ZMC9-1 (F20-T263)
Gene ID:	284266
Molecular Weight:	30-40 kDa

### PROPERTIES

AA Sequence	<pre> F V R T K I D T T E   N L L N T E V H S S   P A Q R W S M Q V P   P E V S A E A G D A A V L P C T F T H P   H R H Y D G P L T A   I W R A G E P Y A G   P Q V F R C A A A R G S E L C Q T A L S   L H G R F R L L G N   P R R N D L S L R V   E R L A L A D D R R Y F C R V E F A G D   V H D R Y E S R H G   V R L H V T A A P R   I V N I S V L P S P A H A F R A L C T A   E G E P P P A L A W   S G P A L G N S L A   A V R S P R E G H G H L V T A E L P A L   T H D G R Y T C T A   A N S L G R S E A S   V Y L F R F H G A S G A S T           </pre>
Biological Activity	Immobilized Human Siglec-15-His at 2 µg/mL (100 µl/well) can bind Anti-Human Siglec15 mAb-mFC, The ED <sub>50</sub> is <18.5 µg/mL.
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, 150 mM NaCl, 0.3% CHAPS, 5% Trehalose, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	The Siglec-15 Protein plays a crucial role in cellular interactions by selectively binding to sialylated glycoproteins, indicating a specific affinity for molecules with sialic acid residues. Additionally, Siglec-15 engages in molecular associations with TYROBP and HCST, suggesting its involvement in intricate signaling pathways. This ability to interact with key signaling
------------	---

---

partners underscores Siglec-15's potential significance in mediating immune responses and cellular communication. The specific recognition of sialylated glycoproteins highlights the protein's role in recognizing and responding to cell surface modifications, contributing to the complex network of cellular interactions.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA