

## IL-16 Protein, Mouse (His)

<b>Cat. No.:</b>	HY-P72590
<b>Synonyms:</b>	Pro-interleukin-16; Interleukin-16; IL-16; Lymphocyte chemoattractant factor; LCF
<b>Species:</b>	Mouse
<b>Source:</b>	E.coli
<b>Accession:</b>	O54824 (S1205-S1322)
<b>Gene ID:</b>	16170
<b>Molecular Weight:</b>	14-16 kDa

### PROPERTIES

<b>AA Sequence</b>	<p>S A A S A S A A S D    I S V E S K E A T V    C T V T L E K T S A    G L G F S L E G G K</p> <p>G S L H G D K P L T    I N R I F K G T E Q    G E M V Q P G D E I    L Q L A G T A V Q G</p> <p>L T R F E A W N V I    K A L P D G P V T I    V I R R T S L Q C K    Q T T A S A D S</p>
<b>Biological Activity</b>	Data is not available.
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20 mM Tris, 150 mM NaCl, pH 8.0.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
<b>Storage &amp; Stability</b>	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA