

FGF-21 Protein, Mouse (HEK293, His)

Cat. No.:	HY-P72651
Synonyms:	FGF21; Fibroblast Growth Factor 21; FGF-21
Species:	Mouse
Source:	HEK293
Accession:	Q9JJN1 (A29-S210)
Gene ID:	56636
Molecular Weight:	20-29 kDa

PROPERTIES

AA Sequence	<p> A Y P I P D S S P L L Q F G G Q V R Q R Y L Y T D D D Q D T E A H L E I R E D G T V V G A A H R S P E S L L E L K A L K P G V I Q I L G V K A S R F L C Q Q P D G A L Y G S P H F D P E A C S F R E L L L E D G Y N V Y Q S E A H G L P L R L P Q K D S P N Q D A T S W G P V R F L P M P G L L H E P Q D Q A G F L P P E P P D V G S S D P L S M V E P L Q G R S P S Y A S </p>
Biological Activity	Measured in a cell proliferation assay using NIH-3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 0.9543 µg/mL in the presence of 1.25 µg/mL Recombinant Human Klotho beta, corresponding to a specific activity is 1047.88 units/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 100 mM NaCl, pH 9.0 or 20 mM PB, 150 mM NaCl, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	FGF-21 protein exerts its biological influence by promoting glucose uptake in differentiated adipocytes, primarily through inducing the expression of the glucose transporter SLC2A1/GLUT1, rather than SLC2A4/GLUT4. This activity is likely contingent on the presence of KLB. Beyond its localized effects, FGF-21 plays a crucial role in the regulation of systemic
------------	--

glucose homeostasis and insulin sensitivity. The protein interacts directly with KLB, facilitated by its C-terminus, and also engages with FGFR4, indicating a complex interplay in its molecular mechanisms.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA