

CEACAM5 Protein, Human (HEK293, Fc)

Cat. No.:	HY-P72693
Synonyms:	Carcinoembryonic antigen-related cell adhesion molecule 5; CEA; CD66e; CEACAM5
Species:	Human
Source:	HEK293
Accession:	P06731 (K35-A685)
Gene ID:	1048
Molecular Weight:	155-180 kDa

PROPERTIES

AA Sequence

K L T I E S T P F N	V A E G K E V L L L	V H N L P Q H L F G	Y S W Y K G E R V D
G N R Q I I G Y V I	G T Q Q A T P G P A	Y S G R E I I Y P N	A S L L I Q N I I Q
N D T G F Y T L H V	I K S D L V N E E A	T G Q F R V Y P E L	P K P S I S S N N S
K P V E D K D A V A	F T C E P E T Q D A	T Y L W W V N N Q S	L P V S P R L Q L S
N G N R T L T L F N	V T R N D T A S Y K	C E T Q N P V S A R	R S D S V I L N V L
Y G P D A P T I S P	L N T S Y R S G E N	L N L S C H A A S N	P P A Q Y S W F V N
G T F Q Q S T Q E L	F I P N I T V N N S	G S Y T C Q A H N S	D T G L N R T T V T
T I T V Y A E P P K	P F I T S N N S N P	V E D E D A V A L T	C E P E I Q N T T Y
L W W V N N Q S L P	V S P R L Q L S N D	N R T L T L L S V T	R N D V G P Y E C G
I Q N K L S V D H S	D P V I L N V L Y G	P D D P T I S P S Y	T Y Y R P G V N L S
L S C H A A S N P P	A Q Y S W L I D G N	I Q Q H T Q E L F I	S N I T E K N S G L
Y T C Q A N N S A S	G H S R T T V K T I	T V S A E L P K P S	I S S N N S K P V E
D K D A V A F T C E	P E A Q N T T Y L W	W V N G Q S L P V S	P R L Q L S N G N R
T L T L F N V T R N	D A R A Y V C G I Q	N S V S A N R S D P	V T L D V L Y G P D
T P I I S P P D S S	Y L S G A N L N L S	C H S A S N P S P Q	Y S W R I N G I P Q
Q H T Q V L F I A K	I T P N N N G T Y A	C F V S N L A T G R	N N S I V K S I T V
S A S G T S P G L S	A		

Appearance Lyophilized powder.

Formulation Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.2.

Endotoxin Level <1 EU/µg, determined by LAL method.

Reconstitution It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH₂O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).

Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

CEACAM5 protein, a cell surface glycoprotein, assumes a multifaceted role in cell adhesion, intracellular signaling, and tumor progression. Functioning as a mediator of both homophilic and heterophilic cell adhesion, CEACAM5 interacts with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM6. In the context of tumor progression, CEACAM5 acts as an oncogene, promoting tumor advancement and inducing resistance to anoikis in colorectal carcinoma cells. Additionally, during microbial infection, CEACAM5 serves as a receptor for E. coli Dr adhesins, and the binding of these adhesins results in the dissociation of the CEACAM5 homodimer. These diverse functions underscore the versatility of CEACAM5 in regulating cellular processes and highlight its significance in both physiological and pathological contexts.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA