

CD79B Protein, Rhesus Macaque (HEK293, His)

Cat. No.:	HY-P72708
Synonyms:	B-cell antigen receptor complex-associated protein beta chain; CD79b; B29; IGB
Species:	Rhesus Macaque
Source:	HEK293
Accession:	H9ZFT8 (A30-D161)
Gene ID:	718218
Molecular Weight:	20-30 kDa

PROPERTIES

AA Sequence	<p>A K S E D L Y P N P K G S A C S R I W Q S P R F I A R K R G F T V K M H C Y V T</p> <p>N S T F S I V S W L R K R E T D K E P Q Q V N L E Q G H M H Q T Q N S S V T T L</p> <p>I I Q D I R F E D N G I Y F C Q Q E C S K T S E V Y R G C G T E L R V M G F S T</p> <p>L A Q L K Q R N T L K D</p>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>CD79B Protein plays an essential role in conjunction with CD79A in initiating the signal transduction cascade triggered by the B-cell antigen receptor complex (BCR). This pivotal function leads to the internalization of the complex, subsequent trafficking to late endosomes, and eventual antigen presentation. CD79B enhances the phosphorylation of CD79A, potentially by recruiting kinases that phosphorylate CD79A or by engaging proteins that bind to CD79A, safeguarding it from dephosphorylation. Forming a heterodimer with the alpha chain, CD79B forms a disulfide-linked complex. It is an integral component of the B-cell antigen receptor complex, where the alpha/beta chain heterodimer associates non-covalently with an antigen-specific membrane-bound surface immunoglobulin comprising two heavy chains and two light chains. Additionally, CD79B interacts with LYN, further contributing to its regulatory role in B-cell signaling.</p>
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA