

CD55/DAF Protein, Human (HEK293, His)

Cat. No.:	HY-P72721
Synonyms:	Complement Decay-Accelerating factor; CD55; CR; DAF
Species:	Human
Source:	HEK293
Accession:	P08174 (D35-S353)
Gene ID:	1604
Molecular Weight:	50-75 kDa

PROPERTIES

AA Sequence	<pre> DCGLPPDVPN AQPALLEGRTS FPEDTVITYK CEESFVKIPG EKDSVICLKG SQWSDIEEFC NRSCEVPTL NSASLKQPYI TQNYFPVGTV VEYECRPGYR REPSLSPKLT CLQNLKWSTA VEFCCKKSCP NPGEIRNGQI DVPGGILFGA TISFSCNTGY KLVFGSTSSFC LISGSSVQWS DPLPECREIY CPAPPQIDNG IIQGERDHYG YRQSVTYACN KGFTMIGEHS IYCTVNNDG EWSGPPPECR GKS L TSKVPP TVQKPTTVNV PTTEVSPTSQ KTTTKTTTPN AQA TRSTPVS RTTKHFHETT PNKGS GTTS </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	CD55/DAF Protein plays a crucial role in the immune system by recognizing C4b and C3b fragments generated during C4 and C3 activation. Its interaction with cell-associated C4b and C3b polypeptides interferes with their ability to catalyze the conversion of C2 and factor B to enzymatically active C2a and Bb, preventing the formation of C4b2a and C3bBb, the amplification convertases of the complement cascade. This interference serves as a regulatory mechanism, inhibiting
------------	---

complement activation and preventing the formation of C3 and C5 convertases, thereby mitigating complement-induced damage. Notably, CD55/DAF also acts as a receptor for Coxsackievirus A21, as well as coxsackieviruses B1, B3, and B5 during microbial infection, highlighting its diverse roles in immune regulation and viral recognition.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA