

Inhibitors • Screening Libraries •

Proteins

CSF1R Protein, Mouse (HEK293, Fc)

Macrophage colony-stimulating factor 1 receptor; CSF-1R; M-CSF-R; CD115; CSF1R; FMS			

PROPERTIES

AA Sequence					
	MELGPPLVLL	LATVWHGQGA	PVIEPSGPEL	VVEPGETVTL	
	R C V S N G S V E W	DGPISPYWTL	D P E S P G S T L T	Т В А Т	
	TYRCTELEDP	MAGSTTIHLY	VKDPAHSWNL	LAQEVTVVEG	
	QEAVLPCLIT	DPALKDSVSL	MREGGRQVLR	KTVYFFSPWR	
	GFIIRKAKVL	D S N T Y V C K T M	VNGRESTSTG	IWLKVNRVHP	
	EPPQIKLEPS	KLVRIRGEAA	QIVCSATNAE	VGFNVILKRG	
	DTKLEIPLNS	DFQDNYYKKV	RALSLNAVDF	QDAGIYSCVA	
	SNDVGTRTAT	MNFQVVESAY	LNLTSEQSLL	QEVSVGDSLI	
	LTVHADAYPS	IQHYNWTYLG	PFFEDQRKLE	FITQRAIYRY	
	TFKLFLNRVK	ASEAGQYFLM	A Q N K A G W N N L	TFELTLRYPP	
	EVSVTWMPVN	GSDVLFCDVS	GYPQPSVTWM	ECRGHTDRCD	
	EAQALQVWND	ТНРЕVLSQКР	FDKVIIQSQL	РІСТЬКНИМТ	
	YFCKTHNSVG	NSSQYFRAVS	LGQSKQLPDE	S	
Appearance	Lyophilized powder.				
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are				
Tornitiation	added as protectants before lyophilization.				
Endotoxin Level	<1 EU/µg, determined by LAL method.				
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH_2O.				
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is				
otoruge a otasinty	recommended to freeze aliquots at -20°C or -80°C for extended storage.				
Shipping	Room temperature in continental US; may vary elsewhere.				

DESCRIPTION

Background

CSF1R protein, a tyrosine-protein kinase, serves as a cell-surface receptor for CSF1 and IL34, exerting pivotal control over the survival, proliferation, and differentiation of hematopoietic precursor cells, particularly mononuclear phagocytes like macrophages and monocytes. Its role in innate immunity and inflammatory processes is underscored by its promotion of the release of pro-inflammatory chemokines in response to IL34 and CSF1. Additionally, CSF1R plays a critical role in the regulation of osteoclast proliferation and differentiation, bone resorption, and is indispensable for normal bone and tooth development. It is essential for normal fertility in both males and females, as well as for the development of milk ducts and acinar structures in the mammary gland during pregnancy. Notably, CSF1R influences the reorganization of the actin cytoskeleton, regulates the formation of membrane ruffles, cell adhesion, cell migration, and facilitates cancer cell invasion. Upon ligand binding, CSF1R activates multiple signaling pathways, including ERK1/2, JNK, PI3K/AKT, MAP kinases (MAPK1/ERK2 and/or MAPK3/ERK1), and SRC family kinases (SRC, FYN, YES1). Its downstream effects involve the phosphorylation of various target proteins, such as PIK3R1, PLCG2, GRB2, SLA2, and CBL. Furthermore, CSF1R mediates the activation of STAT family members (STAT3, STAT5A, and/or STAT5B) and promotes tyrosine phosphorylation of SHC1 and INPP5D/SHIP-1. The receptor signaling is tightly regulated by protein phosphatases, including INPP5D/SHIP-1, and by the rapid internalization of the activated receptor. In the central nervous system, CSF1R may contribute to the development of microglia macrophages.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA