

Animal-Free FGF-2 Protein, Mouse (His)

Cat. No.:	HY-P73052AF
Synonyms:	rMubFGF; HBGF-2; FGF-2; FGF-b; FGF-basic
Species:	Mouse
Source:	E. coli
Accession:	P15655 (A11-S154)
Gene ID:	14173
Molecular Weight:	Approximately 17.2 kDa

PROPERTIES

AA Sequence	<div> <div>A L P E D G G A A F</div> <div>V R E K S D P H V K</div> <div>L A S K C V T E E C</div> <div>Q Y K L G S K T G P</div> </div> <div> <div>P P G H F K D P K R</div> <div>L Q L Q A E E R G V</div> <div>F F F E R L E S N N</div> <div>G Q K A I L F L P M</div> </div> <div> <div>L Y C K N G G F F L</div> <div>V S I K G V C A N R</div> <div>Y N T Y R S R K Y S</div> <div>S A K S</div> </div> <div> <div>R I H P D G R V D G</div> <div>Y L A M K E D G R L</div> <div>S W Y V A L K R T G</div> </div>
Biological Activity	Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is <1.5 ng/mL. The specific activity of recombinant mouse FGF-2 is approximately >1x10 ⁶ IU/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a solution containing 0.01% sarkosyl in 1X PBS, pH 8.0.
Endotoxin Level	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	FGF-2/bFGF is a member of the fibroblast family and has a high affinity for heparin. FGF-2 plays an important role in tendon to bone healing, cartilage repair, bone repair, and nerve regeneration. FGF-2 specifically binds to tyrosine kinase receptors and activates the FGF/FGFR signaling pathway. Subsequently, FGF-2 influences cell proliferation, differentiation and apoptosis, as well as immune regulation by transducing other classical pathways. For example, FGF-2 regulates the JAK-STAT signaling pathway to regulate cartilage metabolism. FGF-2 also acts as a mitotic promoter to accelerate cell proliferation. Therefore, (1) FGF-2 is an important growth factor in the healing process of ligament/tendon injury. In vitro
------------	--

experiments, low-dose FGF-2 can stimulate the proliferation and differentiation of bone marrow mesenchymal stem cells, and up-regulate the mRNA expression of type I/III collagen and fibronectin. However, high doses of FGF-2 did not stimulate extracellular matrix (ECM) protein proliferation and gene expression. (2) FGF-2 is also an endogenous and intrinsic growth factor in cartilage repair. FGF-2 binds to heparan sulfate proteoglycan and is stored in the ECM of articular cartilage. When cartilage is damaged or degenerated, ECM rapidly releases FGF-2 and activates ERK signaling pathways to promote cartilage regeneration. FGF-2 exhibits a biphasic effect in combination with its specific receptor. FGF-2 combined with FGFR3 promoted the repair of articular cartilage. FGF-2 combined with FGFR1 promoted the degeneration of articular cartilage^[1]. FGF-2 is expressed in granulosa cells and colliculus cells, as well as hepatocellular cancer cells, but not in non-cancerous liver tissues. This reveals the role of FGF-2 in brain tumors, particularly glioblastoma. According to studies, FGF-2 is a known carcinogenic factor in GBM. FGF-2 increases the self-renewal of glioblastoma stem cells and contributes to the growth and vascularization of glioma^[2]. FGF-2 protein is highly conserved in some species, and the similarity rate of human FGF-2 protein sequence to rat, mouse, and bovine was 97.4%, 95.45%, and 98.71%, respectively.

REFERENCES

- [1]. Zhang J, et al. FGF2: a key regulator augmenting tendon-to-bone healing and cartilage repair. *Regen Med*. 2020 Sep;15(9):2129-2142.
- [2]. Jimenez-Pascual A, et al. FGF2: a novel druggable target for glioblastoma? *Expert Opin Ther Targets*. 2020 Apr;24(4):311-318.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA