

## IL-10 Protein, Pig

Cat. No.:	HY-P73154
Synonyms:	Interleukin-10 receptor subunit beta; IL-10RB; CRF2-4; IL-10R2; CDw210b
Species:	Pig
Source:	E.coli
Accession:	Q29055 (S19-N175)
Gene ID:	397106
Molecular Weight:	Approximately 18.0 kDa

### PROPERTIES

<b>Biological Activity</b>	Measured by its binding ability in a functional ELISA. Immobilized IL-10 Protein, Pig at 2 µg/mL (100 µl/well) can bind Rhesus IL10RA-hFc and the EC <sub>50</sub> is 25-100 ng/mL. Measured in a cell proliferation assay using MC/9-2 mouse mast cells and the ED <sub>50</sub> is typically 1-5 ng/mL.
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
<b>Storage &amp; Stability</b>	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA