

Noggin Protein, Mouse (HEK293, Fc)

Cat. No.:	HY-P73323
Synonyms:	NOG; Noggin; SYM1; SYNS1; SYNS1A
Species:	Mouse
Source:	HEK293
Accession:	P97466 (Q28-C232)
Gene ID:	18121
Molecular Weight:	55-65 kDa

PROPERTIES

AA Sequence	<p>Q H Y L H I R P A P S D N L P L V D L I E H P D P I F D P K E K D L N E T L L R</p> <p>S L L G G H Y D P G F M A T S P P E D R P G G G G G P A G G A E D L A E L D Q L</p> <p>L R Q R P S G A M P S E I K G L E F S E G L A Q G K K Q R L S K K L R R K L Q M</p> <p>W L W S Q T F C P V L Y A W N D L G S R F W P R Y V K V G S C F S K R S C S V P</p> <p>E G M V C K P S K S V H L T V L R W R C Q R R G G Q R C G W I P I Q Y P I I S E</p> <p>C K C S C</p>
Biological Activity	Measured by its ability to inhibit BMP-4-induced alkaline phosphatase production by ATDC5 mouse chondrogenic cells. The ED ₅₀ for this effect is 0.04-0.3 µg/mL in the presence of 50 ng/mL of recombinant Human BMP4.
Appearance	Lyophilized powder.
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	Noggin is a vital protein involved in cartilage morphogenesis and joint formation, playing a crucial role in inhibiting bone morphogenetic proteins (BMP) signaling essential for the growth and patterning of the neural tube and somite. Acting as an inhibitor of BMP, Noggin is implicated in the regulation of chondrocyte differentiation through its interaction with growth
------------	---

and differentiation factor 5 (GDF5) and likely GDF6. Existing as a homodimer, Noggin directly interacts with GDF5, leading to the inhibition of chondrocyte differentiation. These multifaceted functions underscore the importance of Noggin in embryonic development, particularly in the intricate processes of skeletal and neural tissue formation, and highlight its role as a key modulator of BMP signaling pathways (

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA