

## TrkC Protein, Human (HEK293, His)

Cat. No.:	HY-P73465
Synonyms:	TrkC; NT-3 Growth Factor Receptor; NTRK3; TRKC
Species:	Human
Source:	HEK293
Accession:	Q16288 (M1-D428)
Gene ID:	4916
Molecular Weight:	80-90 kDa

### PROPERTIES

Biological Activity	Imobilized human Trkc-His at 10 µg/mL (100 µl/well )can bind biotinylated human NT3, The EC <sub>50</sub> of biotinylated human NT3 is 0.03-0.07 µg/mL.
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.2 µm filtered solution of Lyophilized from sterile PBS, pH 7.4, 5% Trehalose, 5% Mannitol, 0.01% Tween-80.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	TrkC Protein, a receptor tyrosine kinase, plays a crucial role in nervous system and potential heart development. Upon binding to its ligand NTF3/neurotrophin-3, TrkC undergoes autophosphorylation, initiating various signaling pathways such as the phosphatidylinositol 3-kinase/AKT and the MAPK pathways. These pathways are instrumental in regulating cell survival and differentiation, highlighting the pivotal role of TrkC in orchestrating essential cellular processes during development.
------------	---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA