

## Product Data Sheet

## TOP1 Protein, Human (765a.a, sf9, His)

Cat. No.:	HY-P73584
Synonyms:	DNA topoisomerase 1; TOP1
Species:	Human
Source:	Sf9 insect cells
Accession:	P11387 (M1-F765)
Gene ID:	7150
Molecular Weight:	Approximately 93.1 kDa

PROPERTIES         Biological Activity         The enzyme activity of this recombinant protein is testing in progress, we cannot offer a guarantee yet.
<b>Biological Activity</b> The enzyme activity of this recombinant protein is testing in progress, we cannot offer a guarantee yet.
Appearance         Solution.
<b>Formulation</b> Supplied as a 0.2 μm filtered solution of 20 mM Tris, pH 7.5, 200 mM NaCl, 20% Glycerol.
<b>Endotoxin Level</b> <1 EU/µg, determined by LAL method.
Reconsititution N/A.
<b>Storage &amp; Stability</b> Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping Shipping with dry ice

## DESCRIPTION

Background TOP1 protein plays a pivotal role in cellular processes by alleviating the supercoiling and torsional tension of DNA generated during DNA replication and transcription. This is achieved through the transient cleavage and subsequent rejoining of one strand of the DNA duplex. The enzyme introduces a single-strand break via transesterification at a specific target site in duplex DNA, leading to the formation of a DNA-(3'-phosphotyrosyl)-enzyme intermediate. This intermediate expels a 5'-OH DNA strand, allowing the free DNA strand to rotate around the intact phosphodiester bond on the opposing strand, effectively removing DNA supercoils. In the final religation step, the DNA 5'-OH attacks the covalent intermediate, leading to the expulsion of the active-site tyrosine and restoration of alternative splicing, exemplified by its impact on tissue factor (F3) pre-mRNA in endothelial cells. Additionally, it contributes to the circadian transcription of the core circadian clock component BMAL1 by modifying the chromatin structure surrounding the ROR response elements (ROREs) on the BMAL1 promoter.

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA