

# **Screening Libraries**

Proteins



# Siglec-3/CD33 Protein, Mouse (HEK293)

Cat. No.: HY-P73641

Myeloid Cell Surface Antigen CD33; Siglec-3; gp67; CD33; SIGLEC3 Synonyms:

Species: HEK293 Source:

Accession: Q63994 (M1-E240)

Gene ID: 12489

Molecular Weight: Approximately 25.6 kDa

# **PROPERTIES**

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

# **DESCRIPTION**

Background

Siglec-3/CD33, a sialic-acid-binding immunoglobulin-like lectin, plays a crucial role in mediating cell-cell interactions and maintaining immune cells in a resting state. It exhibits a preference for binding sialic acid on the short O-linked glycans of specific mucins. The protein forms homodimers through disulfide linkages and interacts with signaling molecules such as PTPN6/SHP-1 and PTPN11/SHP-2 upon phosphorylation. Additionally, CD33 engages with C1QA via its C-terminus, leading to the activation of CD33 inhibitory motifs.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1