

S100A8-S100A9 Heterodimer Protein, Human (sf9, Flag-His)

Cat. No.:	HY-P73670
Synonyms:	CAGA; CAGB; CFAG; MRP14; MRP8&S100A9 Protein; S100A8; S100A9
Species:	Human
Source:	Sf9 insect cells
Accession:	P05109 (M1-E93)&P06702 (M1-P114)
Gene ID:	6279&6280
Molecular Weight:	Approximately 12.2&14.2 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris, 500 mM NaCl, 10% Glycerol, pH 8.0. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>S100A8, plays a crucial role in regulating inflammatory processes and immune responses. Often found as calprotectin (S100A8/A9), it serves diverse intracellular functions, including facilitating leukocyte arachidonic acid trafficking and metabolism, modulating the tubulin-dependent cytoskeleton during phagocyte migration, and activating the neutrophilic NADPH-oxidase. In particular, it activates NADPH-oxidase by aiding in the assembly of the enzyme complex at the cell membrane, transferring arachidonic acid, and directly binding to NCF2/P67PHOX. Extracellularly, it exhibits pro-inflammatory, antimicrobial, oxidant-scavenging, and apoptosis-inducing activities. Acting as an alarmin or danger-associated molecular pattern (DAMP) molecule, S100A8 stimulates innate immune cells through binding to pattern recognition receptors such as Toll-like receptor 4 (TLR4) and receptor for advanced glycation endproducts (AGER), activating MAP-kinase and NF-kappa-B signaling pathways and amplifying the pro-inflammatory cascade. With antimicrobial activity against bacteria and fungi, it likely exerts this effect through Zn(2+) chelation essential for microbial growth. Additionally, S100A8/A9 induces cell death via autophagy and apoptosis, regulates neutrophil number and apoptosis, and acts as an oxidant scavenger to prevent tissue damage. It acts as an amplifier of inflammation in autoimmunity and cancer development, and in microbial infection, such as by SARS-CoV-2, it may induce expansion of aberrant immature neutrophils in a TLR4-dependent manner.</p>
-------------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA