

PAM Protein, Human (HEK293, Fc)

Cat. No.:	HY-P73732
Synonyms:	Peptidyl-glycine alpha-amidating monooxygenase; PAM; PHM; PAL
Species:	Human
Source:	HEK293
Accession:	P19021-2 (M1-V710)
Gene ID:	5066
Molecular Weight:	Approximately 104 kDa

PROPERTIES

Biological Activity	The enzyme activity of this recombinant protein is testing in progress, we cannot offer a guarantee yet.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The PAM protein operates as a bifunctional enzyme, overseeing the post-translational modification of inactive peptidylglycine precursors to their bioactive alpha-amidated peptide forms—a crucial terminal modification in the biosynthesis of numerous neural and endocrine peptides. The alpha-amidation process involves two sequential reactions, each governed by distinct catalytic domains within the enzyme. In the first step, the peptidyl alpha-hydroxylating monooxygenase (PHM) domain catalyzes a copper-, ascorbate-, and O₂-dependent stereospecific hydroxylation (with S stereochemistry) at the alpha-carbon (C-alpha) of the C-terminal glycine of the peptidylglycine substrate. The subsequent step, orchestrated by the peptidylglycine amidoglycolate lyase (PAL) domain, entails a zinc-dependent cleavage of the N-C-alpha bond, resulting in the production of the alpha-amidated peptide and glyoxylate. Additionally, PAM exhibits a similar capacity to catalyze the two-step conversion of an N-fatty acylglycine to a primary fatty acid amide and glyoxylate in a manner reminiscent of its peptidylglycine modification function.</p>
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA