

IL-8/CXCL8 Protein, Human (CHO)

Cat. No.:	HY-P7379
Synonyms:	rHuIL-8/CXCL8; C-X-C motif chemokine 8; NAP-1; IL8; GCP-1
Species:	Human
Source:	CHO
Accession:	P10145 (A23-S99)
Gene ID:	3576
Molecular Weight:	Approximately 12.5 kDa

PROPERTIES

AA Sequence	A V L P R S A K E L R C Q C I K T Y S K P F H P K F I K E L R V I E S G P H C A N T E I I V K L S D G R E L C L D P K E N W V Q R V V E K F L K R A E N S
Biological Activity	1. The ED ₅₀ is <10 ng/mL as measured by CHO/Gα15 cells. 2. Measured in a cell proliferation assay using HUVEC human umbilical vein endothelial cells. The ED ₅₀ for this effect is 1.656 ng/mL, corresponding to a specific activity is 6.039×10 ⁵ units/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS or 20 mM PB, 150 mM NaCl, pH 7.4.
Endotoxin Level	<0.2 EU/μg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>IL-8 (CXCL8) belongs to the ELR⁺ CXC chemokines family. IL-8 is initially produced as a protein of 99 amino acids that undergoes cleavage to form active IL-8 isoforms, a 77 amino acid peptide in non-immune cells or a 72 amino acid peptide in monocytes and macrophages. The gene encoding IL-8 is located on chromosome 4q13-q21. Dimerisation of IL-8 forms the structural basis for receptor binding. IL-8 is expressed by various cells including monocytes, macrophages, leukocytes, endothelial cells, and epithelial cells^{[1][2][3]}.</p> <p>Mature human IL-8/CXCL8 shares 75% amino acid sequence identity with canine IL-8/CXCL8. While, human IL-8 shares</p>
------------	--

94.95% aa sequence identity with Rhesus Macaque IL-8 protein.

IL-8 is responsible for the recruitment and activation of neutrophils and granulocytes to the site of inflammation. IL-8 is almost undetectable in physiological states, but is rapidly induced by pro-inflammatory cytokines such as TNF α and IL-1 β . The function of IL-8 mainly relies on its interaction with specific cell surface GPCR, CXCR1 and CXCR2. In addition, IL-8 is reported to promote integrin β 3 upregulation and the invasion of hepatocellular carcinoma cells through activation of the PI3K/Akt pathway. In odontogenic lesions, IL-8 has been proven to be highly expressed in ameloblastoma epithelial cells and irreversible pulpitis. In rheumatoid arthritis and other inflammatory joint diseases IL-8 could bring about the accumulation of neutrophils, which are considered a major source of cartilage-degrading enzymes. IL-8 stimulates the MAPK and tyrosine phosphorylation of cellular proteins. Tumour cells and fibroblasts communicate with each other, including autocrine and paracrine factors, including IL-8, resulting in the upregulation of MMP2 and MMP9 degradable extracellular matrix (ECM) components that trigger tumour invasion^{[1][2][3][4]}.

IL-8 is typically known to promote angiogenesis, but it also activates matrix metalloproteinase (MMP) that is involved in metastasis-related tissue remodelling. IL-8 is induced in lipopolysaccharide (LPS)-stimulated monocytes and shown to induce neutrophil migration. IL-8 exerts multiple effects on biological activities of tumour cells including proliferation, invasion and migration. IL-8 also increases the expression of Akt in androgen-independent prostate cancer (AIPC) cell lines. IL-8 activates MAPK signalling via PI3K in neutrophils, and via transactivation of EGFR resulting in Ras-GTPase activation in ovarian and lung cancer cell lines. There is substantial amount of experimental data suggesting that IL-8 and receptors contribute to elimination of pathogens, but may also contribute significantly to disease-associated processes, including tissue injury, fibrosis, angiogenesis and tumorigenesis^{[3][5]}.

REFERENCES

- [1]. M Baggiolini, et al. Neutrophil-activating peptide-1/interleukin 8, a novel cytokine that activates neutrophils. *J Clin Invest.* 1989 Oct;84(4):1045-9.
- [2]. M Wolf, et al. Granulocyte chemotactic protein 2 acts via both IL-8 receptors, CXCR1 and CXCR2. *Eur J Immunol.* 1998 Jan;28(1):164-70.
- [3]. Qian Liu, et al. The CXCL8-CXCR1/2 pathways in cancer. *Cytokine Growth Factor Rev.* 2016 Oct;31:61-71.
- [4]. Jian-Feng Liu, et al. IL-8 Is Upregulated in the Tissue-Derived EVs of Odontogenic Keratocysts. *Biomed Res Int.* 2022 Jul 30;2022:9453270.
- [5]. Remo C Russo, et al. The CXCL8/IL-8 chemokine family and its receptors in inflammatory diseases. *Expert Rev Clin Immunol.* 2014 May;10(5):593-619.
- [6]. Hai Jiang, et al. CXCL8 promotes the invasion of human osteosarcoma cells by regulation of PI3K/Akt signaling pathway. *APMIS.* 2017 Sep;125(9):773-780.
- [7]. L Laterveer, et al. Rapid mobilization of hematopoietic progenitor cells in rhesus monkeys by a single intravenous injection of interleukin-8. *Blood.* 1996 Jan 15;87(2):781-8.
- [8]. Baggiolini M, et al. Neutrophil-activating peptide-1/interleukin 8, a novel cytokine that activates neutrophils. *J Clin Invest.* 1989 Oct;84(4):1045-9.
- [9]. Koch AE, et al. Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science.* 1992 Dec 11;258(5089):1798-801.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA