

IL-8/CXCL8 Protein, Human (72a.a, HEK293, His)

Cat. No.:	HY-P73874
Synonyms:	C-X-C motif chemokine 8; MDNCF; Emoctakin; NAP-1
Species:	Human
Source:	HEK293
Accession:	P10145 (S28-S99)
Gene ID:	3576
Molecular Weight:	11-13 kDa

PROPERTIES

Biological Activity	Immobilized Human IL-8, His Tag at 0.5 µg/mL (100µl/well) on the plate. Dose response curve for Anti-IL-8 Antibody, hFc Tag with the EC ₅₀ of <13.6 ng/mL determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 8% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

IL-8 (CXCL8) belongs to the ELR⁺ CXC chemokines family. IL-8 is initially produced as a protein of 99 amino acids that undergoes cleavage to form active IL-8 isoforms, a 77 amino acid peptide in non-immune cells or a 72 amino acid peptide in monocytes and macrophages. The gene encoding IL-8 is located on chromosome 4q13-q21. Dimerisation of IL-8 forms the structural basis for receptor binding. IL-8 is expressed by various cells including monocytes, macrophages, leukocytes, endothelial cells, and epithelial cells^{[1][2][3]}.

Mature human IL-8/CXCL8 shares 75% amino acid sequence identity with canine IL-8/CXCL8. While, human IL-8 shares 94.95% aa sequence identity with Rhesus Macaque IL-8 protein.

IL-8 is responsible for the recruitment and activation of neutrophils and granulocytes to the site of inflammation. IL-8 is almost undetectable in physiological states, but is rapidly induced by pro-inflammatory cytokines such as TNFα and IL-1β. The function of IL-8 mainly relies on its interaction with specific cell surface GPCR, CXCR1 and CXCR2. In addition, IL-8 is reported to promote integrin β3 upregulation and the invasion of hepatocellular carcinoma cells through activation of the PI3K/Akt pathway. In odontogenic lesions, IL-8 has been proven to be highly expressed in ameloblastoma epithelial cells

and irreversible pulpitis. In rheumatoid arthritis and other inflammatory joint diseases IL-8 could bring about the accumulation of neutrophils, which are considered a major source of cartilage-degrading enzymes. IL-8 stimulates the MAPK and tyrosine phosphorylation of cellular proteins. Tumour cells and fibroblasts communicate with each other, including autocrine and paracrine factors, including IL-8, resulting in the upregulation of MMP2 and MMP9 degradable extracellular matrix (ECM) components that trigger tumour invasion^{[1][2][3][4]}.

IL-8 is typically known to promote angiogenesis, but it also activates matrix metalloproteinase (MMP) that is involved in metastasis-related tissue remodelling. IL-8 is induced in lipopolysaccharide (LPS)-stimulated monocytes and shown to induce neutrophil migration. IL-8 exerts multiple effects on biological activities of tumour cells including proliferation, invasion and migration. IL-8 also increases the expression of Akt in androgen-independent prostate cancer (AIPC) cell lines. IL-8 activates MAPK signalling via PI3K in neutrophils, and via transactivation of EGFR resulting in Ras-GTPase activation in ovarian and lung cancer cell lines. There is substantial amount of experimental data suggesting that IL-8 and receptors contribute to elimination of pathogens, but may also contribute significantly to disease-associated processes, including tissue injury, fibrosis, angiogenesis and tumorigenesis^{[3][5]}.

REFERENCES

- [1]. M Baggiolini, et al. Neutrophil-activating peptide-1/interleukin 8, a novel cytokine that activates neutrophils. *J Clin Invest*. 1989 Oct;84(4):1045-9.
 - [2]. M Wolf, et al. Granulocyte chemotactic protein 2 acts via both IL-8 receptors, CXCR1 and CXCR2. *Eur J Immunol*. 1998 Jan;28(1):164-70.
 - [3]. Qian Liu, et al. The CXCL8-CXCR1/2 pathways in cancer. *Cytokine Growth Factor Rev*. 2016 Oct;31:61-71.
 - [4]. Jian-Feng Liu, et al. IL-8 Is Upregulated in the Tissue-Derived EVs of Odontogenic Keratocysts. *Biomed Res Int*. 2022 Jul 30;2022:9453270.
 - [5]. Remo C Russo, et al. The CXCL8/IL-8 chemokine family and its receptors in inflammatory diseases. *Expert Rev Clin Immunol*. 2014 May;10(5):593-619.
 - [6]. Hai Jiang, et al. CXCL8 promotes the invasion of human osteosarcoma cells by regulation of PI3K/Akt signaling pathway. *APMIS*. 2017 Sep;125(9):773-780.
 - [7]. L Laterveer, et al. Rapid mobilization of hematopoietic progenitor cells in rhesus monkeys by a single intravenous injection of interleukin-8. *Blood*. 1996 Jan 15;87(2):781-8.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA