

# **Screening Libraries**

**Proteins** 



## **Product** Data Sheet

# **B7-H4 Protein, Human (HEK293, His)**

Cat. No.: HY-P74395

Synonyms: V-set domain containing T-cell activation inhibitor 1; VTCN1; Protein B7S1; B7-H4

Species: HEK293 Source:

Accession: Q7Z7D3 (F29-A258)

Gene ID: 79679 **Molecular Weight:** 43-48 kDa

**PROPERTIES** 

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/μg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH2O.

recommended to freeze aliquots at -20°C or -80°C for extended storage.

Room temperature in continental US; may vary elsewhere.

### **DESCRIPTION**

Storage & Stability

Background

Shipping

B7-H4 protein functions as a negative regulator of the T-cell-mediated immune response, exerting inhibitory effects on Tcell activation, proliferation, cytokine production, and the development of cytotoxicity. Particularly significant when expressed on the cell surface of tumor macrophages, B7-H4, in collaboration with regulatory T-cells (Treg), plays a crucial role in suppressing tumor-associated antigen-specific T-cell immunity. Additionally, B7-H4 is implicated in the promotion of epithelial cell transformation, indicating its multifaceted involvement in modulating immune responses and cellular processes associated with tumor development.

Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1