

TNFRSF13B Protein, Human (HEK293, His)

Cat. No.:	HY-P74506
Synonyms:	Tumor necrosis factor receptor superfamily member 13B; TACI; CD267; Tnfrsf13b
Species:	Human
Source:	HEK293
Accession:	O14836-2 (S2-T120)
Gene ID:	23495
Molecular Weight:	18-24 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	TNFRSF13B Protein serves as the receptor for TNFSF13/APRIL and TNFSF13B/TALL1/BAFF/BLYS, demonstrating high-affinity binding to both ligands. Its activation results in calcineurin-dependent activation of NF-AT, along with the activation of NF-kappa-B and AP-1, thereby playing a crucial role in stimulating B- and T-cell function and regulating humoral immunity. Additionally, TNFRSF13B binds TRAF2, TRAF5, and TRAF6, suggesting its involvement in various signaling pathways. Notably, it interacts with the NH2-terminal domain of CAMLG using its C-terminus.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA