

## IFN-alpha 1/IFNA1 Protein, Mouse (P.pastoris, His)

Cat. No.:	HY-P74866
Synonyms:	IFNA1; IFNalpha 1; IFN-alpha 1
Species:	Mouse
Source:	P. pastoris
Accession:	P01572 (C24-K189)
Gene ID:	15962
Molecular Weight:	Approximately 20.5 kDa

### PROPERTIES

Biological Activity	Measured in antiviral assays using L929 cells infected with vesicular stomatitisvirus (VSV) and the ED <sub>50</sub> is 20-100 pg/mL.
Appearance	Solution.
Formulation	Supplied as a 0.22 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

### DESCRIPTION

#### Background

IFN-alpha 1 (IFNA1; IFN-α1), belongs to the alpha/beta interferon family, is produced by macrophages with antiviral activities<sup>[1]</sup>. Interferon (IFN) is originally identified as a substance 'interfering' with viral replication in vitro. IFN-α/β and related molecules are classified as type I IFNs, as for the other two types of type II IFN (IFN-γ) and type III IFNs (IFN-λ), respectively<sup>[2]</sup>.

IFNs binds to one of three type-specific receptors, which leads to the activation of JAK1 and TYK2<sup>[3]</sup>. This signal transduction results in phosphorylation of STAT1 and STAT2 and eventually in an association with IFN regulatory factor 9 (IRF9) and formation of the IFN-stimulated gene factor 3 (ISGF3) complex. Thus the ISGF3 complex induces transcription of IFN-stimulated genes (ISGs), with subsequent immunomodulatory effects on both innate and adaptive immune responses<sup>[4]</sup>.

The interactions of type I IFN with the immune system is important for the generation of a durable antitumor response through its effects on dendritic cells (DC)<sup>[5]</sup>. IFN has been widely used for animal disease model, and the sequence of amino acids in IFNA1 protein of mouse is very different from human (62.96%).

### REFERENCES

- 
- [1]. Zoon KC, et al. Purification and characterization of multiple components of human lymphoblastoid interferon-alpha. J Biol Chem. 1992 Jul 25;267(21):15210-6.
- [2]. Zhang SY, et al. Inborn errors of interferon (IFN)-mediated immunity in humans: insights into the respective roles of IFN-alpha/beta, IFN-gamma, and IFN-lambda in host defense. Immunol Rev. 2008 Dec;226:29-40.
- [3]. Gibbert K, et al. IFN- $\alpha$  subtypes: distinct biological activities in anti-viral therapy. Br J Pharmacol. 2013 Mar;168(5):1048-58.
- [4]. De Ceuninck F, et al. IFN- $\alpha$ : A key therapeutic target for multiple autoimmune rheumatic diseases. Drug Discov Today. 2021 Oct;26(10):2465-2473.
- [5]. Lapenta C, et al. IFN-Alpha-Mediated Differentiation of Dendritic Cells for Cancer Immunotherapy: Advances and Perspectives. Vaccines (Basel). 2020 Oct 19;8(4):617.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA