

## GITR Protein, Mouse (HEK293, Fc)

Cat. No.:	HY-P75165
Synonyms:	Tumor necrosis factor receptor superfamily member 18; CD357; TNFRSF18; AITR; GITR
Species:	Mouse
Source:	HEK293
Accession:	O35714 (M1-H153)
Gene ID:	21936
Molecular Weight:	Approximately 41 kDa

### PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	<p>GITR is expressed on regulatory T cells (Tregs) and some activated immune cells, including effector T lymphocytes, nature killer (NK) cells, and neutrophils<sup>[1]</sup>.</p> <p>The amino acid sequence of human GITR protein has low homology for mouse GITR protein.</p> <p>GITR does not have any enzymatic activity and signaling is propagated via recruiting TRAF-family members, specifically TRAF1, TRAF2 and TRAF5, to the GITR-signaling complex. The signaling is then mediated through NF-κB and MAPK pathways. GITR does not have any enzymatic activity and signaling is propagated via recruiting TRAF-family members, specifically TRAF1, TRAF2 and TRAF5, to the GITR-signaling complex. The signaling is then mediated through NF-κB and MAPK pathways, protecting T cells from TCR activation-induced cell death<sup>[2]</sup>.</p> <p>GITR (Glucocorticoid-induced TNFR-related protein, also known as TNFRSF18) is a type I transmembrane protein. GITR stimulates the proliferation of effector T-lymphocytes and partially reverses the immunosuppressive function of CD4+CD25+ Tregs<sup>[1]</sup>. GITR is activated by its ligand GITRL (TNFSF18). GITR induces NOS in murine macrophage in a time and dose-dependent manner<sup>[3]</sup>. GITR inhibits Multiple Myeloma (MM) cell proliferation in vitro and in vivo and induces apoptosis<sup>[4]</sup>.</p>
------------	--

### REFERENCES

- 
- [1]. Tian J, et al. The Role of GITR/GITRL Interaction in Autoimmune Diseases. Front Immunol. 2020 Oct 9;11:588682.
- [2]. Krausz LT, et al. GITR-GITRL system, a novel player in shock and inflammation. ScientificWorldJournal. 2007 May 1;7:533-66.
- [3]. Shin HH, et al. Recombinant glucocorticoid induced tumor necrosis factor receptor (rGITR) induces NOS in murine macrophage. FEBS Lett. 2002 Mar 13;514(2-3):275-80.
- [4]. Liu Y, et al. Novel tumor suppressor function of glucocorticoid-induced TNF receptor GITR in multiple myeloma. PLoS One. 2013 Jun 13;8(6):e66982.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA