

## CD96 Protein, Human (Biotinylated, HEK293, His)

Cat. No.:	HY-P75371
Synonyms:	T-cell surface protein tactile; Cell surface antigen CD96; CD96
Species:	Human
Source:	HEK293
Accession:	P40200-2 (V22-M503)
Gene ID:	10225
Molecular Weight:	120-130 kDa

### PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/ $\mu$ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	CD96 protein appears to play a role in adhesive interactions during the late phase of the immune response, potentially involving activated T and NK cells. It facilitates NK cell-target adhesion by interacting with PVR present on target cells. CD96 may function in a phase after T and NK cells have penetrated the endothelium using integrins and selectins, actively engaging with diseased cells and navigating within areas of inflammation. Structurally, CD96 forms homodimers through disulfide linkages and interacts specifically with PVR, suggesting its involvement in cell-cell interactions crucial for immune responses during inflammation and infection.
------------	---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA