

# **Screening Libraries**

## **Product** Data Sheet

### CD83 Protein, Rat (HEK293, His)

Cat. No.: HY-P75375

Synonyms: B-cell activation protein; CD83 antigen; hCD83; CD83; BL11

Species:

HEK293 Source:

B2GV95 (M22-A134) Accession:

Gene ID: 361226

Molecular Weight: Approximately 27-35 kDa due to the glycosylation.

#### **PROPERTIES**

	C		
$\Delta \Delta$	Sec	ша	nco

MREVTVACSE TADLPCTAPW DPQLSYTVSW AKVSESGNER LELPESKQNS SVEAPKKRPY SLTIQNTTIC SAGTYRCALQ ELGGQRNFSG TVVLKVTGCP

KEATESTFRK YRA

**Biological Activity** 

Measured by the ability of the immobilized protein to support the adhesion of human monocyte-derived dendritic cells. When 5 x 10<sup>4</sup> cells/well are added to Rat CD83 coated plates (5 μg/mL, 100 μL/well), approximately 68.76% adhered after 30 minutes at 37° C.

**Appearance** 

Lyophilized powder

**Formulation** 

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

**Endotoxin Level** 

<1 EU/µg, determined by LAL method.

Reconsititution

It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH<sub>2</sub>O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).

Storage & Stability

Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping

Room temperature in continental US; may vary elsewhere.

#### **DESCRIPTION**

Background

CD83 Protein is suggested to play a significant role, potentially contributing to antigen presentation or mediating cellular interactions that ensue following lymphocyte activation. The specific mechanisms through which CD83 influences these processes, particularly in the context of immune responses, remain to be fully elucidated. As a monomer, CD83 may participate in distinct molecular events that impact antigen presentation and cellular interactions, highlighting its potential as a key regulatory element in modulating immune responses. In-depth investigations are needed to unravel the intricacies

Page 1 of 2 www.MedChemExpress.com

**Proteins** 

of CD83's function and its implications in the orchestration of immune processes.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com