

CD30/TNFRSF8 Protein, Human (HEK293, hFc)

Cat. No.:	HY-P75421
Synonyms:	CD30L receptor; Tumor necrosis factor receptor superfamily member 8; Ki-1 antigen
Species:	Human
Source:	HEK293
Accession:	P28908 (F19-K379)
Gene ID:	943
Molecular Weight:	Approximately 65.2 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	CD30/TNFRSF8, a receptor for TNFSF8/CD30L, is implicated in the regulation of cellular growth and the transformation of activated lymphoblasts. This receptor plays a role in modulating gene expression by activating NF-kappa-B, a key transcription factor associated with diverse cellular processes. The interaction of CD30/TNFRSF8 with signaling adapters such as TRAF1, TRAF2, TRAF3, and TRAF5 underscores its involvement in intricate cellular signaling networks. This receptor's engagement with TNFSF8 suggests its potential impact on immune responses and cellular homeostasis, highlighting its significance in the regulation of fundamental biological processes.
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA