

BMP-5 Protein, Human (HEK293, hFc)

Cat. No.:	HY-P75472
Synonyms:	Bone morphogenetic protein 5; BMP-5
Species:	Human
Source:	HEK293
Accession:	NP_066551.1 (Q324-H454)
Gene ID:	653
Molecular Weight:	50-55 kDa

PROPERTIES

Biological Activity	Measured by its ability to induce alkaline phosphatase production by ATDC5 mouse chondrogenic cells. The ED ₅₀ for this effect is 0.7568 µg/mL, corresponding to a specific activity is 1.32×10 ³ U/mg.
Appearance	Solution
Formulation	Supplied as a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice

DESCRIPTION

Background

Bone Morphogenetic Protein 5 (BMP-5) is a ligand protein with pleiotropic, belongs to TGFβ family and expresses mainly in the lung and liver. BMP/TGFβ signaling to involve in vascular and valvular homeostasis, which is a critical process of embryonic development^[4]. And BMP/TGFβ signaling can be terminated by inhibitory SMADs including SMAD6 and SMAD7, which are activated and induced by BMP signaling and switch off BMP signaling via multiple mechanisms^[5]. BMP-5 silencing inhibits chondrocyte senescence and apoptosis as well as osteoarthritis (OA) progression by downregulating activity in the p38/ERK signaling pathway^[1]. BMP-5 initiates the canonical BMP signaling cascade by associating with type I receptor BMPRI1A and type II receptor BMPRII^[2]. BMP-5 also triggers signal through non-canonical pathway such as MAPK p38 signaling cascade to promote chondrogenic differentiation^[3]. BMP-5 specific expressing in neural crest progenitor cells is sufficient to induce cell proliferation through the MEK-ERK-ID3 signaling cascade, whereas disruption of this signaling cascade had no effect on cell survival^[6].

REFERENCES

- [1]. Shao Y, et al. BMP5 silencing inhibits chondrocyte senescence and apoptosis as well as osteoarthritis progression in mice. *Aging (Albany NY)*. 2021 Mar 19;13(7):9646-9664.
- [2]. Beck HN, et al. Bone morphogenetic protein-5 (BMP-5) promotes dendritic growth in cultured sympathetic neurons. *BMC Neurosci*. 2001;2:12.
- [3]. Snelling SJ, et al. BMP5 activates multiple signaling pathways and promotes chondrogenic differentiation in the ATDC5 growth plate model. *Growth Factors*. 2010 Aug;28(4):268-79.
- [4]. Yang P, et al. The role of bone morphogenetic protein signaling in vascular calcification. *Bone*. 2020 Dec;141:115542.
- [5]. Miyazawa K, et al. Regulation of TGF- β Family Signaling by Inhibitory Smads. *Cold Spring Harb Perspect Biol*. 2017 Mar 1;9(3):a022095.
- [6]. Shih HY, et al. Bmp5 Regulates Neural Crest Cell Survival and Proliferation via Two Different Signaling Pathways. *Stem Cells*. 2017 Apr;35(4):1003-1014.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA