

B7-2/CD86 Protein, Human (239a.a, HEK293, His)

Cat. No.:	HY-P75491
Synonyms:	Activation B7-2 antigen; B70; BU63; CD86; CD28LG2
Species:	Human
Source:	HEK 293
Accession:	P42081-3 (M1-H239)
Gene ID:	942
Molecular Weight:	55-60 kDa

PROPERTIES

Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized human CD86 at 20 µg/mL (100 µL/well) can bind human CD28 with a linear range of 32-800 ng/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA