

Fc gamma RIIA/CD32a Protein, Human (HEK293, Fc)

Cat. No.:	HY-P75644
Synonyms:	Low Affinity Immunoglobulin Gamma Fc Region Receptor II-a; CD32; FCGR2A; FCG2; IGFR2
Species:	Human
Source:	HEK 293
Accession:	P12318 (M1-I218)
Gene ID:	2212
Molecular Weight:	Approximately 55 kDa

PROPERTIES

Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized human CD32a-Fc at 10 µg/mL (100 µL/well) can bind biotinylated human IgG1 with a linear range of 0.625-10 µg/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized a 0.2 µm filtered solution of PBS, pH 7.5.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA