

## Siglec-3/CD33 Protein, Cynomolgus/Rhesus Macaque (Biotinylated, HEK293, His)

Cat. No.:	HY-P75652
Synonyms:	Myeloid Cell Surface Antigen CD33; Siglec-3; gp67; CD33; SIGLEC3
Species:	Rhesus Macaque
Source:	HEK293
Accession:	A0A2K5W2R5 (M1-G248)
Gene ID:	102117580
Molecular Weight:	Approximately 27.3 kDa

### PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	Siglec-3/CD33, a sialic-acid-binding immunoglobulin-like lectin, plays a crucial role in mediating cell-cell interactions and maintaining immune cells in a resting state. It exhibits a preference for binding sialic acid on the short O-linked glycans of specific mucins. The protein forms homodimers through disulfide linkages and interacts with signaling molecules such as PTPN6/SHP-1 and PTPN11/SHP-2 upon phosphorylation. Additionally, CD33 engages with C1QA via its C-terminus, leading to the activation of CD33 inhibitory motifs.
------------	---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA