

CD79B Protein, Rat (HEK293, Fc)

Cat. No.:	HY-P75658
Synonyms:	B-cell antigen receptor complex-associated protein beta chain; CD79b; B29; IGB
Species:	Rat
Source:	HEK293
Accession:	O70153 (V26-D158)
Gene ID:	171055
Molecular Weight:	Approximately 50 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The human, murine, and rat B29 (Ig beta, CD79b) genes are highly conserved in sequence and organization and exhibit strict B cell-specific expression. In the human and rat genomes, the B29 gene is located between the skeletal muscle-specific Na-channel alpha subunit (SCN4A) gene and the pituitary-specific growth hormone (GH-N) gene. Rat B29/Ig-β gene was 3.1 kb in length with six exons and was separated by 3.3 and 9.3 kb from Na-channel and GH genes, respectively^{[1][2]}.</p> <p>The B-cell receptor (BCR) consists of surface-bound immunoglobulin (Ig) and a heterodimeric signaling unit comprised of CD79A and CD79B. Upon cognate antigen recognition, the receptor initiates important signals for B-cell development and function^[3].</p>
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA