

Fractalkine/CX3CL1 Protein, Canine (HEK293)

Cat. No.:	HY-P75777
Synonyms:	Fractalkine; CX3CL1; FKN; SCYD1; Neurotactin; NTT
Species:	Canine
Source:	HEK293
Accession:	H1ADY9 (Q25-R379)
Gene ID:	487265
Molecular Weight:	64-69 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	Fractalkine/CX3CL1 protein is a member of the intercrine beta (chemokine CC) family, positioning it within a category of chemokines with pivotal roles in intercellular communication and immune responses. As part of the intercrine beta family, Fractalkine/CX3CL1 likely plays a significant role in modulating inflammatory processes and cellular interactions. Further exploration is essential to uncover the specific functions and implications of Fractalkine/CX3CL1 within the broader framework of the chemokine CC family, shedding light on its importance in mediating immune responses and contributing to the intricate network of intercellular signaling.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA