

## BCA-1/CXCL13 Protein, Cynomolgus (His)

Cat. No.:	HY-P76164
Synonyms:	C-X-C motif chemokine 13; BCA1; BLC; SCYB13
Species:	Cynomolgus
Source:	E. coli
Accession:	A0A2K5V497 (V23-P109)
Gene ID:	/
Molecular Weight:	Approximately 14 kDa.

### PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 10 mM Tris, 250 mM NaCl, pH 8.0. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

#### Background

CXCL13, also known as B lymphocyte chemoattractant, is originally identified in stromal cells in B cell follicles as regulating homing of B cells and subsets of T cells. CXCL13 plays a key role in orchestrating cell migration within spatially distinct regions of the secondary lymphoid organs. It strongly attracts B lymphocytes while promoting migration of only small numbers of T cells and macrophages. CXCL13 and its receptor, CXCR5, play fundamental roles in inflammatory, infectious, cancer and immune responses<sup>[1][2][3]</sup>.

The amino acid sequence of human CXCL13 protein has low homology with mouse CXCL13 protein.

CXCL13 exerts its functions through its receptor CXCR5. CXCR5 is highly expressed on mature recirculating B-lymphocytes, a subpopulation of follicular helper T cells (TFH) and skin-derived migratory dendritic cells (DCs), and controls their migration into secondary lymphoid organs towards the gradient of CXCL13. As the loss of the BLR1/CXCR5 receptor is sufficient to disrupt organization of follicles in spleen and Peyer's patches, BCA-1 may act as a B cell homing chemokine. Human BCA-1 competes with radiolabeled IFN-γ inducible protein 10 (IP-10) for binding to the human CXCR3 receptor expressed in Ba/F3 and 293EBNA cell lines. Furthermore, human BCA-1 is an efficacious attractant for human CXCR3 transfected cells. BCA-1 does not induce calcium release in B-lymphocytes. In addition, human BCA-1 is an agonist in stimulating GTP gamma S binding. Human BCA-1 is a specific and functional G-protein-linked chemotactic ligand for the human CXCR3 receptor.

CXCL13 has been widely implicated in the pathogenesis of a number of autoimmune diseases and inflammatory conditions,

---

as well as in lymphoproliferative disorders. In addition, the CXCL13:CXCR5 axis orchestrates cell-cell interactions that regulate lymphocyte infiltration within the tumor microenvironment<sup>[1][2][3]</sup>.

Dysregulation of the CXCL13:CXCR5 axis affecting both B- and TFH cell function is major player in autoimmune disorders, and potentially serves as a biomarker for disease progression and therapeutic response. Moreover, expression of CXCR5 and CXCL13 is shown to be dysregulated in HIV infection, such that the number of CXCR5+ B cells decreases with progression of HIV infection, together with an increase in plasma levels of CXCL13. CXCL13/CXCR5 signaling modulates cancer cell ability to grow, proliferate, invade, and metastasize. CXCL13 drives spinal astrocyte activation and neuropathic pain via CXCR5<sup>[1][2][3]</sup> [4].

---

## REFERENCES

- [1]. Jenh CH, et al. Human B cell-attracting chemokine 1 (BCA-1; CXCL13) is an agonist for the human CXCR3 receptor. *Cytokine*. 2001 Aug 7;15(3):113-21.
- [2]. Muzammal Hussain, et al. CXCL13/CXCR5 signaling axis in cancer. *Life Sci*. 2019 Jun 15;227:175-186.
- [3]. Marcelo G Kazanietz, et al. CXCL13 and Its Receptor CXCR5 in Cancer: Inflammation, Immune Response, and Beyond. *Front Endocrinol (Lausanne)*. 2019 Jul 12;10:471.
- [4]. Bao-Chun Jiang, et al. CXCL13 drives spinal astrocyte activation and neuropathic pain via CXCR5. *J Clin Invest*. 2016 Feb;126(2):745-61.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA