

Product Data Sheet

Screening Libraries

Proteins

CD200 Protein, Rhesus Macaque (HEK293, Fc)

Cat. No.: HY-P76211

OX-2 membrane glycoprotein; CD200; MOX1; MOX2; My033 Synonyms:

Species: Rhesus Macaque

HEK293 Source:

Accession: H9EXH3 (Q31-G232)

Gene ID: 708110

Molecular Weight: Approximately 49.6 kDa.

			IES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/μg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

The CD200 protein plays a pivotal role in immune modulation, functioning as a costimulatory molecule that promotes T-cell proliferation. Additionally, CD200 is implicated in the potential regulation of myeloid cell activity across various tissues, as suggested by similarity-based inferences. The interaction between CD200 and its receptor, CD200R1, is facilitated through their respective N-terminal Ig-like domains. This interaction likely contributes to the regulatory processes involved in immune responses and myeloid cell function. The engagement of CD200 with CD200R1 underscores its significance in mediating immune cell communication and homeostasis, emphasizing its potential as a key player in immunomodulation.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1