

DNAM-1 Protein, Rat (HEK293, Fc)

Cat. No.:	HY-P76214
Synonyms:	CD226 antigen; DNAX accessory molecule 1; DNAM-1; CD226
Species:	Rat
Source:	HEK293
Accession:	D3ZS97 (M1-I265)
Gene ID:	307199
Molecular Weight:	62-68 kDa.

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/ μ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	The DNAM-1 protein is intricately involved in intercellular adhesion, lymphocyte signaling, cytotoxicity, and lymphokine secretion mediated by cytotoxic T-lymphocytes (CTL) and natural killer (NK) cells. Serving as a cell surface receptor for NECTIN2, DNAM-1, upon ligand binding, stimulates T-cell proliferation and the production of cytokines such as IL2, IL5, IL10, IL13, and IFNG. Furthermore, DNAM-1 competes with PVRIG for NECTIN2-binding, underscoring its regulatory role in cellular interactions and signaling pathways. This protein also interacts with PVR and NECTIN2, contributing to its multifaceted functions in immune responses and cellular communication.
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA