

# **Screening Libraries**

**Proteins** 



## **Product** Data Sheet

# CD83 Protein, Cynomolgus (HEK293, Fc)

Cat. No.: HY-P76245

B-cell activation protein; CD83 antigen; hCD83; CD83; BL11 Synonyms:

Species: Cynomolgus HEK293 Source:

Accession: F6WX37 (M1-A143)

Gene ID: 701825

Molecular Weight: 45-55&34 kDa.

Р	Кί	JP	ĿК	Ш	ES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### **DESCRIPTION**

Background

CD83 Protein is suggested to play a significant role, potentially contributing to antigen presentation or mediating cellular interactions that ensue following lymphocyte activation. The specific mechanisms through which CD83 influences these processes, particularly in the context of immune responses, remain to be fully elucidated. As a monomer, CD83 may participate in distinct molecular events that impact antigen presentation and cellular interactions, highlighting its potential as a key regulatory element in modulating immune responses. In-depth investigations are needed to unravel the intricacies of CD83's function and its implications in the orchestration of immune processes.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1