

CD83 Protein, Cynomolgus (HEK293, His)

Cat. No.:	HY-P76246
Synonyms:	B-cell activation protein; CD83 antigen; hCD83; CD83; BL11
Species:	Cynomolgus
Source:	HEK293
Accession:	F6WX37 (T20-A143)
Gene ID:	701825
Molecular Weight:	Approximately 19&21&27 kDa.

PROPERTIES

AA Sequence	<p>T P E V K V A C S E D V D L P C T A P W D P Q V P Y T V S W V K L L E G G E E R</p> <p>M E T P Q E D H L R G Q H Y H Q K G Q N G S L D A P S E R P Y S L K I R N T T S</p> <p>C N S G A Y R C T L Q D P D G Q R N L S G K V I L R V T G C P A Q R K E E T F K</p> <p>K Y R A</p>
Biological Activity	Measured by the ability of the immobilized protein to support the adhesion of human monocyte-derived dendritic cells. When 5x10 ⁴ cells/well are added to Cynomolgus CD83 coated plates (5 µg/mL, 100 µL/well), approximately 65.15% adhered after 30 min at 37°C.
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	CD83 Protein is suggested to play a significant role, potentially contributing to antigen presentation or mediating cellular interactions that ensue following lymphocyte activation. The specific mechanisms through which CD83 influences these processes, particularly in the context of immune responses, remain to be fully elucidated. As a monomer, CD83 may participate in distinct molecular events that impact antigen presentation and cellular interactions, highlighting its potential
------------	---

as a key regulatory element in modulating immune responses. In-depth investigations are needed to unravel the intricacies of CD83's function and its implications in the orchestration of immune processes.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA