

CRIP2 Protein, Human (sf9, His-GST)

Cat. No.:	HY-P76291
Synonyms:	Cysteine-rich protein 2; CRP-2; Protein ESP1
Species:	Human
Source:	Sf9 insect cells
Accession:	P52943 (M1-P208)
Gene ID:	1397
Molecular Weight:	Approximately 49 kDa.

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20 mM Tris, 500 mM NaCl, pH 7.4, 10% Glycerol. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/ μ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	CRIP2 protein, also known as Cysteine-rich intestinal protein 2, is a protein that is involved in various cellular processes. CRIP2 protein interacts with TGFB111. TGFB111, also known as Transforming Growth Factor Beta 1 Induced Transcript 1, is a protein that is induced by transforming growth factor beta 1 (TGF- β 1) and is involved in TGF- β signaling pathways. The interaction between CRIP2 and TGFB111 suggests a potential role for CRIP2 in modulating TGF- β signaling or being involved in TGF- β -mediated cellular processes. However, further research is required to fully understand the functional significance of this interaction and the precise role of CRIP2 protein in cellular mechanisms regulated by TGF- β 1 and TGFB111.
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA