

Product Data Sheet

ST2/IL1RL1 Protein, Mouse (332a.a, HEK293, His)

| Cat. No.: | HY-P76662 |
|-------------------|---|
| Synonyms: | Interleukin-1 receptor-like 1; Interleukin-33 receptor alpha chain; Il1rl1; Ly84; St2; Ste2 |
| Species: | Mouse |
| Source: | HEK293 |
| Accession: | P14719 (M1-R332) |
| Gene ID: | 17082 |
| Molecular Weight: | Approximately 55.7 kDa |

| PROPERTIES | |
|----------------------------|--|
| | |
| Biological Activity | The enzyme activity of this recombinant protein is testing in progress, we cannot offer a guarantee yet. |
| Appearance | Lyophilized powder. |
| Formulation | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. |
| Endotoxin Level | <1 EU/µg, determined by LAL method. |
| Reconsititution | It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH_2O. |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage. |
| Shipping | Room temperature in continental US; may vary elsewhere. |

| DESCRIPTION | |
|-------------|--|
| Background | ST2/IL1RL1 serves as a receptor for interleukin-33 (IL-33), playing crucial roles in innate and adaptive immunity to maintain tissue homeostasis and respond to environmental stresses, in conjunction with its coreceptor IL1RAP. Upon stimulation, it recruits MYD88, IRAK1, IRAK4, and TRAF6, leading to the phosphorylation of various MAPKs, including MAPK3/ERK1, MAPK1/ERK2, MAPK14, and MAPK8. This receptor is implicated in helper T-cell function and, in the context of tissue injury, induces a UCP2-dependent mitochondrial rewiring. This rewiring attenuates the generation of reactive oxygen species, preserving the integrity of the Krebs cycle. This preservation is essential for sustained production of itaconate and subsequent GATA3-dependent differentiation of inflammation-resolving alternatively activated macrophages. Additionally, ST2/IL1RL1 plays a role in inhibiting IL-33 signaling. |

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA