

CD3 epsilon Protein, Rhesus Macaque (HEK293, Fc)

Cat. No.:	HY-P76789
Synonyms:	T-Cell Surface Glycoprotein CD3 Epsilon Chain; T-Cell Surface Antigen T3/Leu-4 Epsilon Chain; CD3e; CD3E; T3E
Species:	Rhesus Macaque
Source:	HEK293
Accession:	XP_001097204 (M1-D117)
Gene ID:	699467
Molecular Weight:	Approximately 38 kDa.

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	The CD3 epsilon protein, a vital component of the TCR-CD3 complex on T-lymphocytes, is pivotal for adaptive immune responses. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. CD3E is crucial for proper T-cell development and contributes to TCR-CD3 complex internalization and down-regulation. The CD3D/CD3E and CD3G/CD3E heterodimers form trimers with TCRalpha and TCRbeta, completing the TCR-CD3 complex. CD3E's interactions with CD6 and NCK1 highlight its multifaceted role in T-cell responses ^[1] .
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA