

CNPY2 Protein, Human (HEK293, His)

Cat. No.:	HY-P76839
Synonyms:	Protein canopy homolog 2; MSAP; TMEM4; ZSIG9
Species:	Human
Source:	HEK293
Accession:	Q9Y2B0-1 (R21-S178)
Gene ID:	10330
Molecular Weight:	Approximately 19 kDa

PROPERTIES

AA Sequence	<pre> RRSQDLHCGA CRALVDELEW EIAQVDPKKT IQMGSFRINP DGSQSVVEVP YARSEAHLTE LLEEICDRMK EYGEQIDPST HRKNYVRVVG RNGESSELDL QGIRIDSDIS GTLKFACESEI VEEYEDELIE FFSREADNVK DKLCSKRTDL CDHALHIS </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	CNPY2, a positive regulator of neurite outgrowth, exerts its function by stabilizing the myosin regulatory light chain (MRLC). It plays a crucial role in preventing MIR-mediated ubiquitination of MRLC, thereby averting its subsequent proteasomal degradation. The interaction between CNPY2 and MYLIP/MIR underscores its regulatory influence on MRLC stability, emphasizing its significance in facilitating neurite outgrowth.
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA