

Inhibitors

Proteins

Screening Libraries

Product Data Sheet

Hemagglutinin/HA Protein, H11N9 (ABB87228, HEK293, His)

Cat. No.: HY-P77016

Synonyms: Influenza A H11N9 (A/mallard/Alberta/294/1977) Hemagglutinin / HA Protein (His)

Species: HEK293 Source:

Accession: ABB87228 (M1-K528)

Gene ID:

Molecular Weight: 65-70 kDa

| Ρ | | | | | | | |
|---|-----|---|---|-----|------|---|----|
| | 184 | u | _ | P.4 | | _ | ٠. |

| Appearance | Lyophilized powder. | | |
|---------------------|--|--|--|
| Formulation | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. | | |
| Endotoxin Level | <1 EU/μg, determined by LAL method. | | |
| Reconsititution | It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. | | |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage. | | |
| Shipping | Room temperature in continental US; may vary elsewhere. | | |

DESCRIPTION

Background

The Hemagglutinin (HA) protein is central to the attachment of virus particles to host cells by binding to sialic acidcontaining receptors on the cell surface. This interaction leads to virion internalization through either clathrin-dependent endocytosis or a clathrin- and caveolin-independent pathway. HA plays a pivotal role in determining host range restriction and virulence, functioning as a Class I viral fusion protein responsible for the penetration of the virus into the cell cytoplasm. Its mediation of the fusion between the membrane of the endocytosed virus particle and the endosomal membrane is essential for successful infection. The acidic environment in endosomes induces an irreversible conformational change in the HA2 subunit, releasing the fusion hydrophobic peptide. The formation of a competent fusion pore requires the cooperative action of several HA trimers, highlighting the intricate molecular processes orchestrated by HA during viral entry.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1